



ISPD gene

isoprenoid synthase domain containing

Normal Function

The *ISPD* gene provides instructions for making a protein that is involved in a process called glycosylation. Through this chemical process, sugar molecules are added to certain proteins. In particular, the ISPD protein helps produce a molecule called ribitol 5-phosphate, which is an important component of the chain of sugar molecules added to a protein called alpha (α)-dystroglycan. Glycosylation is critical for the normal function of α -dystroglycan.

The α -dystroglycan protein helps anchor the structural framework inside each cell (cytoskeleton) to the lattice of proteins and other molecules outside the cell (extracellular matrix). In skeletal muscles, glycosylated α -dystroglycan helps stabilize and protect muscle fibers. In the brain, it helps direct the movement (migration) of nerve cells (neurons) during early development.

Health Conditions Related to Genetic Changes

limb-girdle muscular dystrophy

Walker-Warburg syndrome

At least 17 mutations in the *ISPD* gene have been found to cause Walker-Warburg syndrome, the most severe form of a group of disorders known as congenital muscular dystrophies. Walker-Warburg syndrome causes skeletal muscle weakness and abnormalities of the brain and eyes. Because of the severity of the problems caused by this condition, affected individuals usually do not survive past early childhood.


ISPD gene mutations involved in Walker-Warburg syndrome prevent the normal glycosylation of α -dystroglycan. As a result, α -dystroglycan can no longer effectively anchor cells to the proteins and other molecules that surround them. Without functional α -dystroglycan to stabilize the muscle fibers, they become damaged as they repeatedly contract and relax with use. The damaged fibers weaken and die over time, which affects the development, structure, and function of skeletal muscles in people with Walker-Warburg syndrome.

Defective α -dystroglycan also affects the migration of neurons during the early development of the brain. Instead of stopping when they reach their intended destinations, some neurons migrate past the surface of the brain into the fluid-filled space that surrounds it. Researchers believe that this problem with neuronal migration causes a brain abnormality called cobblestone lissencephaly, in which the surface of the brain lacks the normal folds and grooves and instead appears bumpy and irregular. Less is known about the effects of *ISPD* gene mutations on other parts of the body.

Chromosomal Location

Cytogenetic Location: 7p21.2, which is the short (p) arm of chromosome 7 at position 21.2

Molecular Location: base pairs 16,087,527 to 16,422,562 on chromosome 7 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- 2-C-methyl-D-erythritol 4-phosphate cytidylyltransferase-like protein
- 4-diphosphocytidyl-2C-methyl-D-erythritol synthase homolog
- hCG_1745121
- isoprenoid synthase domain-containing protein
- IspD
- ISPD_HUMAN
- MDDGA7
- Nip
- notch1-induced protein

Additional Information & Resources

Educational Resources

- Molecular Cell Biology (fourth edition, 2000): Protein Glycosylation in the ER and Golgi Complex
<https://www.ncbi.nlm.nih.gov/books/NBK21744/>
- Washington University Neuromuscular Disease Center
<http://neuromuscular.wustl.edu/syncm.html#ispd>

GeneReviews

- Congenital Muscular Dystrophy Overview
<https://www.ncbi.nlm.nih.gov/books/NBK1291>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28ISPD%5BTIAB%5D%29+OR+%28IsPD%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- ISOPRENOID SYNTHASE DOMAIN-CONTAINING PROTEIN
<http://omim.org/entry/614631>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_ISPD.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=ISPD%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=37276
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/729920>
- UniProt
<http://www.uniprot.org/uniprot/A4D126>

Sources for This Summary

- Cirak S, Foley AR, Herrmann R, Willer T, Yau S, Stevens E, Torelli S, Brodd L, Kamynina A, Vondracek P, Roper H, Longman C, Korinthenberg R, Marrosu G, Nürnberg P; UK10K Consortium, Michele DE, Plagnol V, Hurles M, Moore SA, Sewry CA, Campbell KP, Voit T, Muntoni F. ISPD gene mutations are a common cause of congenital and limb-girdle muscular dystrophies. *Brain.* 2013 Jan;136(Pt 1):269-81. doi: 10.1093/brain/aws312. Epub 2013 Jan 3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23288328>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3562076/>
- Gerin I, Ury B, Breloy I, Bouchet-Seraphin C, Bolsée J, Halbout M, Graff J, Vertommen D, Muccioli GG, Seta N, Cuisset JM, Dabaj I, Quijano-Roy S, Grahn A, Van Schaftingen E, Bommer GT. ISPD produces CDP-ribitol used by FKTN and FKRP to transfer ribitol phosphate onto α -dystroglycan. *Nat Commun.* 2016 May 19;7:11534. doi: 10.1038/ncomms11534.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/27194101>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4873967/>
- OMIM: ISOPRENOID SYNTHASE DOMAIN-CONTAINING PROTEIN
<http://omim.org/entry/614631>
- Kanagawa M, Kobayashi K, Tajiri M, Manya H, Kuga A, Yamaguchi Y, Akasaka-Manya K, Furukawa J, Mizuno M, Kawakami H, Shinohara Y, Wada Y, Endo T, Toda T. Identification of a Post-translational Modification with Ribitol-Phosphate and Its Defect in Muscular Dystrophy. *Cell Rep.* 2016 Mar 8;14(9):2209-23. doi: 10.1016/j.celrep.2016.02.017.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/26923585>
- Roscioli T, Kamsteeg EJ, Buysse K, Maystadt I, van Reeuwijk J, van den Elzen C, van Beusekom E, Riemersma M, Pfundt R, Vissers LE, Schraders M, Altunoglu U, Buckley MF, Brunner HG, Grisart B, Zhou H, Veltman JA, Gilissen C, Mancini GM, Delrée P, Willemsen MA, Ramadza DP, Chitayat D, Bennett C, Sheridan E, Peeters EA, Tan-Sindhunata GM, de Die-Smulders CE, Devriendt K, Kayserili H, El-Hashash OA, Stemple DL, Lefeber DJ, Lin YY, van Bokhoven H. Mutations in ISPD cause Walker-Warburg syndrome and defective glycosylation of α -dystroglycan. *Nat Genet.* 2012 May;44(5):581-5. doi: 10.1038/ng.2253.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22522421>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3378661/>
- Tasca G, Moro F, Aiello C, Cassandrini D, Fiorillo C, Bertini E, Bruno C, Santorelli FM, Ricci E. Limb-girdle muscular dystrophy with α -dystroglycan deficiency and mutations in the ISPD gene. *Neurology.* 2013 Mar 5;80(10):963-5. doi: 10.1212/WNL.0b013e3182840cbc. Epub 2013 Feb 6.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23390185>
- Willer T, Lee H, Lommel M, Yoshida-Moriguchi T, de Bernabe DB, Venzke D, Cirak S, Schachter H, Vajsar J, Voit T, Muntoni F, Loder AS, Dobyns WB, Winder TL, Strahl S, Mathews KD, Nelson SF, Moore SA, Campbell KP. ISPD loss-of-function mutations disrupt dystroglycan O-mannosylation and cause Walker-Warburg syndrome. *Nat Genet.* 2012 May;44(5):575-80. doi: 10.1038/ng.2252.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22522420>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3371168/>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/ISPD>

Reviewed: January 2017
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services